Osceola (OS) 2020 Rankings

| 0 1 | | 1.7 |
|---------|----------|---------|
| (County | / Demogi | raphics |
| Conne | 20111091 | артпоо |

| | County | State |
|--|--------|-----------|
| Population | 6,040 | 3,156,145 |
| % below 18 years of age | 23.9% | 23.2% |
| % 65 and older | 21.5% | 17.1% |
| % Non-Hispanic Black | 0.5% | 3.8% |
| % American Indian & Alaska Native | 0.6% | 0.5% |
| % Asian | 1.0% | 2.7% |
| % Native Hawaiian/Other Pacific Islander | 0.1% | 0.1% |
| % Hispanic | 8.5% | 6.2% |
| % Non-Hispanic White | 88.3% | 85.3% |
| % not proficient in English | 3% | 2% |
| % Females | 49.0% | 50.2% |
| % Rural | 58.4% | 36.0% |
| | | |
| | | |

| | Osceola County | Error Margin | Top U.S. Performers ^ | Iowa | Rank (of 99) | |
|--|-------------------------|--------------------------------------|--------------------------|-------------------------|--------------|----|
| Health Outcomes | | | | | | 62 |
| Length of Life | | | | | | 87 |
| Premature death Quality of Life | 8,200 | 5,600-11,600 | 5,500 | 6,200 | | 13 |
| Poor or fair health ** Poor physical health days ** Poor mental health days ** Low birthweight | 14% 3.2 3.5 4% | 14-15% 3.0-3.3 3.3-3.7 2-6% | 12% 3.1 3.4 6% | 14% 3.3 3.6 7% | | |

Additional Health Outcomes (not included in overall ranking)

| Life expectancy | 79.1 | 76.6-81.6 | 81.1 | 79.4 |
|----------------------------------|------|-----------|------|------|
| Premature age-adjusted mortality | 350 | 270-440 | 270 | 320 |
| Child mortality | | | 40 | 50 |
| Infant mortality | | | 4 | 5 |
| Frequent physical distress | 9% | 9-10% | 9% | 10% |
| Frequent mental distress | 11% | 11-11% | 11% | 11% |
| Diabetes prevalence | 7% | 3-14% | 7% | 10% |
| HIV prevalence | | | 41 | 98 |
| | | | | |

| Health Factors | | | | |
|------------------------|------|---------|-----|-------|
| Health Behaviors | | | | |
| | | | | |
| Adult smoking ** | 16% | 15-16% | 14% | 17% |
| Adult obesity | 32% | 23-42% | 26% | 33% |
| Food environment index | 8.6 | | 8.6 | 8.2 |
| Dhysical inactivity | 220/ | 15 210/ | 20% | 2.40/ |

| Adult obesity | 32% | 23-42% | 26% | 33% |
|----------------------------------|-------|--------|-------|-------|
| Food environment index | 8.6 | | 8.6 | 8.2 |
| Physical inactivity | 22% | 15-31% | 20% | 24% |
| Access to exercise opportunities | 62% | | 91% | 83% |
| Excessive drinking ** | 19% | 18-20% | 13% | 22% |
| Alcohol-impaired driving deaths | 30% | 13-48% | 11% | 27% |
| Sexually transmitted infections | 148.9 | | 161.4 | 441.6 |
| Teen births | 25 | 17-36 | 13 | 19 |

Additional Health Behaviors (not included in overall ranking)

| Food insecurity | 10% | 9% | 11% |
|---------------------------------|-----|----|-----|
| Limited access to healthy foods | 4% | 2% | 6% |
| Drug overdose deaths | | 10 | 10 |
| Motor vehicle crash deaths | | 9 | 11 |

30

| 21/2020 | Osceola County, | iowa County i i | calli Markings & Moad | шарэ | |
|---------------------------------|--------------------|-------------------|-----------------------|---------|----|
| Insufficient sleep | 27% | 26-28% | 27% | 28% | |
| Clinical Care | | | | | 74 |
| | | | | | |
| Uninsured | 8% | 6-9% | 6% | 5% | |
| Primary care physicians | 3,020:1 | | 1,030:1 | 1,370:1 | |
| Dentists | 6,040:1 | | 1,240:1 | 1,500:1 | |
| Mental health providers | | | 290:1 | 640:1 | |
| Preventable hospital stays | 2,784 | | 2,761 | 3,808 | |
| Mammography screening | 49% | | 50% | 51% | |
| Flu vaccinations | 51% | | 53% | 53% | |
| | | | | | |
| Additional Clinical Care | e (not included in | overall r | anking) | | |
| | • | | J, | | |
| | | | | | |

| Uninsured adults Uninsured children Other primary care providers | 9% 4% 1,510:1 | 7-10% 3-6% | 7% 3% 665:1 | 7% 3% 990:1 | |
|--|---------------------|---------------|-------------------|-------------------|----|
| Social & Economic Factors | | | | | 34 |
| | | | | | |
| High school graduation | 94% | | 96% | 91% | |
| Some college | 55% | 46-64% | 73% | 70% | |
| Unemployment | 1.9% | | 2.6% | 2.5% | |
| Children in poverty | 13% | 9-17% | 11% | 14% | |
| Income inequality | 4.1 | 3.3-5.0 | 3.7 | 4.2 | |
| Children in single-parent households | 23% | 16-30% | 20% | 29% | |
| Social associations | 24.8 | | 18.4 | 15.2 | |
| Violent crime | 90 | | 63 | 282 | |
| Injury deaths | 95 | 64-136 | 58 | 67 | |

Additional Social & Economic Factors (not included in overall ranking)

| Disconnected youth | | | 4% | 5% | |
|---|----------|-----------------|----------|----------|----|
| Reading scores | 3.2 | | 3.4 | 3.1 | |
| Math scores | 3.3 | | 3.4 | 3.1 | |
| Median household income | \$60,600 | \$55,000-66,200 | \$69,000 | \$60,100 | |
| Children eligible for free or reduced price lunch | 44% | | 32% | 40% | |
| Residential segregation - Black/White | | | 23 | 62 | |
| Residential segregation - non-white/white | 9 | | 14 | 48 | |
| Homicides | | | 2 | 2 | |
| Suicides | | | 11 | 14 | |
| Firearm fatalities | | | 8 | 9 | |
| Juvenile arrests | 33 | | | 37 | |
| Physical Environment | | | | | 14 |
| | | | | | |
| Air pollution - particulate matter ** | 8.1 | | 6.1 | 9.0 | |
| Drinking water violations | No | | | | |
| Severe housing problems | 10% | 7-13% | 9% | 12% | |
| Driving alone to work | 80% | 76-84% | 72% | 81% | |
| Long commute - driving alone | 17% | 13-21% | 16% | 20% | |

Additional Physical Environment (not included in overall ranking)

| Traffic volume | 16 | | | 1,461 |
|----------------------------|-----|--------|-----|-------|
| Homeownership | 76% | 73-78% | 81% | 71% |
| Severe housing cost burden | 9% | 5-13% | 7% | 10% |

Areas to Explore

Areas of Strength

Note: Blank values reflect unreliable or missing data

^{^ 10}th/90th percentile, i.e., only 10% are better.

^{**} Data should not be compared with prior years